

## The Association between Surgical Abortion and Preterm Birth: An Overview

*Evidence in peer-reviewed literature from 168 studies over fifty years points to a causal, dose-response relationship between surgical abortion and subsequent preterm birth. This document provides an overview of this literature, discusses mechanisms for this effect, demonstrates the strength of evidence for causality, and offers guidance for informed consent prior to surgical abortion. This document does not provide detailed statistical analysis or a high-resolution assessment of the quality of studies on surgical abortion and preterm birth (covered in Practice Guideline 11).*

### Background

#### *Preterm Birth*

Preterm birth (PTB), defined as birth before 37 weeks of pregnancy, plagues modern society. There are over 3 million annual deaths worldwide due to PTB, and PTB is estimated to cost over 100 million disability-adjusted life-years, when combined with low birth weight (LBW).<sup>1</sup> The incidence of PTB ranges from 6 to 8% in Europe, Australia, and Canada<sup>2-3</sup> to 9 to 12% in Asia, Africa, and is currently 10.1% in the United States, a decrease since the push to eliminate non-indicated PTB.<sup>7, 8</sup>

The literature has shown for some time the increasing risk for PTB with surgical abortion. In 2018, 92% of abortions were before 13 weeks, with about half of them being surgical.<sup>64</sup> Researchers of varying countries and

political bent have found that surgical abortion confers an increased risk for PTB, which may be mediated by infection risk.<sup>32, 34-36</sup>

#### *Evidence for Increased Preterm Birth after Abortion*

As of November 2021, 168 studies have been published on the association between abortion and PTB. A complete review of the literature is provided in Practice Bulletin 11, but this document reviews key studies at a foundational level. The landmark meta-analyses on PTB after abortion are:

- Swingle et al., a 2009 meta-analysis
- Shah et al., a 2009 meta-analysis
- Oppenraajj et al., a 2009 review
- Lowit et al., a 2010 meta-analysis
- Saccone et al., a 2016 meta-analysis

The first landmark study is Swingle et al., which examined studies published between 1995 and 2007 and found that women with a prior abortion had increased odds of delivery before 32 weeks (1.64, 95% CI 1.38-1.91).<sup>44</sup>

A few comments are helpful to understand these results. The increased odds ratio (OR) published by Swingle et al. was 1.64, and it was statistically significant as denoted by the 95% confidence interval (95% CI) of 1.38 to 1.91, which does not include 1.0. A confidence interval denotes 95% certainty that the true difference in odds resides between the two values; if the 95% CI includes 1.0, we cannot be certain that there is no difference from the control group (here, the group with no prior abortion), denoted by their odds of 1.0. Odds are different than relative risk, or absolute risk difference, and require some computation to derive a clinically memorable percent risk. An odds ratio of 1.64 translates to an increase in risk from 1.5% (the United States baseline rate of delivery before 32 weeks) to about 2.4%. Importantly, this is not a 64% increase. That would be reported in a study as a relative risk (RR) of 1.64, different from odds.

The second landmark study from 2009 is Shah et al, which found increased odds of delivery before 37 weeks (OR 1.35, 95% CI 1.20-1.52).<sup>38</sup> These odds mean the rate of birth before 37 weeks after one abortion is 13%, compared to the baseline 10%. This study also reported the odds of PTB after two or more abortions, OR 1.72 (95% CI 1.45-2.04). This translates to an increase in risk

from 10% to about 18%, nearly doubling. Shah et al.'s results also show the important epidemiological principle of a *dose effect*: the more abortions prior to first delivery, the higher the risk for PTB.

Oppenraaij et al. combined 13 studies and found increased risk of delivery before 32 weeks and delivery before 37 weeks after one abortion, and that effect was more dramatic after two or more induced abortions (a dose effect).<sup>45</sup>

Lowit et al. reported data from seven systematic reviews (including four meta-analyses) and eighteen primary studies found increased risk of delivery before 32 weeks and before 37 weeks, concluding that “[c]urrent evidence ... suggest an association between IA [Induced abortion] and pre-term birth.”<sup>46</sup>

Saccone et al. included 36 studies in a systematic review and meta-analysis. This study found that women with one prior abortion had a significantly increased risk of PTB (OR 1.52, 95% CI 1.08-2.16), a significant increase in odds that translates to a risk increase from 10% to 14%.<sup>47</sup>

### *Pathophysiology of Induced Abortion and Preterm Birth*

The putative mechanisms by which surgical induced abortion may increase the risk for PTB may include the following:

1. Cervical trauma from surgical dilation.

2. Predisposition to inflammation, or subclinical inoculation from the procedure.
3. Chronic increased production of maternal stress hormones.

Regarding mechanical trauma, dilation and curettage (D&C) is independently associated with an increased risk of PTB based on the investigation of neutral researchers.<sup>33</sup> The mechanical injury from the surgical procedure itself is the most likely reason that surgical abortion increases PTB risk.<sup>27</sup>

Regarding infection, this hypothesis emerges from the association of infection and inflammation with PTB,<sup>31</sup> coupled with data about the risk of chorioamnionitis during a subsequent delivery. The risk of chorioamnionitis in a pregnancy after abortion is threefold<sup>37</sup> or fourfold<sup>38</sup> higher compared to live birth (OR 4.0, 95% CI 2.7-5.8).

### *Causality in Medicine: Bradford Hill Causation Criteria*

There is substantial evidence for an association between surgical abortion and PTB—more evidence than for the relationship between tobacco use and preterm birth. (This is not to belittle the association between tobacco and PTB, but to show that a neutral observer who acknowledges that association would also acknowledge an abortion-PTB association.)

But before insisting on a response like that to tobacco, we must discuss criteria for determining *causality*, whether one thing is

actually *causing* another, or simply associated with it.

The Bradford Hill criteria have been used since the 1960s for this purpose (see Box 1). Dr. Hill cautioned, however:

I do not believe [there are] hard-and-fast rules ... that must be observed before we accept cause and effect. None of my [criteria is] indisputable evidence for or against the cause-and-effect hypothesis and none can be required as a *sine qua non*. What they can do [is] help us to make up our minds on [whether] there any other answers equally, or more, likely than cause and effect? All scientific work is incomplete [and] liable to be...modified by advancing knowledge. That does not confer ... a freedom to ignore the knowledge we already have, or to postpone ... action.<sup>38</sup>

Thus, while the Bradford Hill criteria are a good foundation, the lack of any particular criterion is not grounds for dismissal of a causal relationship.

### *Applying the Bradford Hill Criteria to Abortion and Preterm Birth*

Here, the comparison between surgical abortion and tobacco use is helpful. In 1964, the US Surgeon General applied the emerging Bradford Hill criteria for causality to studies evaluating the association between tobacco use and PTB, and chose to warn the public of a potential causal effect of tobacco use on risk of PTB.

Box 1. The Bradford Hill Criteria for Causality	
<b>Strength of the association</b>	Does the effect meet statistical and/or clinical significance?
<b>Consistency</b>	Does the effect provide consistent results or outcomes?
<b>Specificity</b>	Is the effect specific to the outcome or result?
<b>Temporality</b>	Does the effect occur prior or during the given item under study?
<b>Dose Response</b>	Does the effect increase with increasing exposure?
<b>Plausibility</b>	Does the effect meet criteria for biologically reasonableness?
<b>Coherence</b>	Does the effect make sense with the outcome specified or found?
<b>Experiment</b>	Is the effect experimentally reproducible in multiple experiments with diverse authors and/or populations?
<b>Analogy</b>	Is the effect similar (analogous) to other effects found experimentally or clinically?
AAPLOG Practice Bulletin 5, Nov 2021.	

With regard to *timing*, surgical abortion occurs before a subsequent pregnancy at risk of PTB. There is a known *dose effect* demonstrated for the risk of PTB and very pre-term (VPTB) birth increasing with a greater number of induced surgical abortions.<sup>31,39</sup> (No such increased risk has been demonstrated with smoking and PTB.)

The *experiment* for surgical abortion has been repeated dozens of times, in over 168 studies on the topic. There is also *consistency* of the effects of prior surgical abortion, and no study shows a protective effect of prior surgical abortion. There is inconsistency on tobacco use and PTB,<sup>40</sup> since some studies show a protective effect of tobacco.<sup>39</sup>

Induced abortion has a very *strong effect* on the rate of subsequent PTB and very preterm birth (delivery before 32 weeks).<sup>32,39</sup> Biologic *plausibility* for prior surgical abortion as a cause for future preterm birth is thought to be the result of either trauma or inflammation mediated, as mentioned above.<sup>29-32</sup> This leads to *coherence* with subsequent evidence of cervical insufficiency or chorioamnionitis. This is *analogous* to the risk of preterm birth from other surgeries that affect cervical integrity (e.g. cervical conization) or on other procedures that may result in intrauterine inflammation.

While the effect of abortion on PTB is not unique (there are other factors that increase risk of PTB), this lack of the criterion of

specificity is common in clinical outcomes. Tobacco is also not the only factor associated with increased risk of PTB, and this non-specificity does not disqualify either tobacco use or surgical abortion as causal in the pathophysiology of PTB.

The logical conclusion drawn from the published literature that linked tobacco use and lung cancer is almost exactly the same as the logical conclusion drawn from the published literature linking induced surgical abortion and PTB: there is a causal relationship.

## Clinical Questions and Answers

Q *This practice bulletin doesn't address some of my concerns about the quality of the evidence available on this purported "link." Who does?*

Practice Bulletin 11 is designed to delve into the quality of evidence available on this link and investigates the statistical and methodological merit of many of the studies on this topic.

Q *What about medication abortion?*

There has not been much data on medication abortion yet, in comparison to the decades of data on surgical abortion. Bhattacharya et al., 2012 found that women with previous abortion (medication or surgical) had increased risk of PTB (adjusted relative risk of 2.3, 95% CI 2.27-2.33). This study had some missing data on tobacco use and type of abortion (not

listed in 25% of cases), which are weaknesses in a study of abortion and PTB.<sup>11</sup>

Q *What do other medical experts say about the relationship between surgical abortion and PTB?*

AAPLOG is the only organization in the United States has formally acknowledged the risk with induced abortion for PTB, but is not alone in its assessment of the evidence.

Dr. Jay Iams is an Associate Editor of the *American Journal of Obstetrics and Gynecology* and editor of a major maternal-fetal medicine textbook. He served as president of the Society for Maternal-Fetal Medicine from 2003-04 and of the American Gynecological and Obstetrical Society in 2013. Dr. Iams is one of the leading researchers in PTB and wrote in 2010,

Contrary to common belief, population-based studies have found that elective pregnancy terminations in the first and second trimester are associated with a very small but apparently real increase in the risk of subsequent spontaneous preterm birth.<sup>41</sup>

Dr. Phil Steer, editor of the *British Journal of Obstetrics and Gynecology* wrote an editorial comment on a major meta-analysis of surgical abortion and PTB,

A key finding is that compared to women with no history of termination, even allowing for the expected higher incidence of socio-economic disadvantage, women with just one [termination of pregnancy] had an increased odds of subsequent

preterm birth. However, finding that even one termination can increase the risk of preterm birth means that we should continue to search for ways of making termination less traumatic.<sup>42</sup>

The Royal College of Obstetrics and Gynaecology (RCOG) acknowledges the association of surgical abortion and PTB. In a 2011 guideline entitled “The Care of Women Requesting Induced Abortion,” RCOG advises:

Women should be informed that induced abortion is associated with a small increase in the risk of subsequent preterm birth, which increases with the number of abortions. However, there is insufficient evidence to imply causality.<sup>43</sup>

Despite 168 peer-reviewed publications documenting an increased risk for PTB with surgical abortion, the leading medical organizations for women’s healthcare including the American College of Obstetricians and Gynecologists (ACOG) refuse to acknowledge the increased associated risk for PTB or acknowledge the substantial body of literature raising this concern, as of their 2016 reaffirmation of Practice Bulletin 130.<sup>25</sup>

Planned Parenthood, the largest provider of abortion in the U.S., does not inform patients of the association of surgical abortion with PTB, instead stating that

[s]afe, uncomplicated abortion does not cause problems for future pregnancies such as birth defects, premature birth or low birth weight babies ...or infant death.<sup>44</sup>

*Q What are the effects of abortion-related preterm birth?*

A conservative estimate for the last 43 (1973-2018) years is approximately 102,056 deaths associated with delivery before 32 weeks related to prior abortion.<sup>23</sup> Of these deaths, 46,268 (45%) are estimated to be of Black infants, an over-representation given that Black Americans represent 15-16% of the total population.<sup>25</sup> As noted by one author, this is “equal to the number of lives...lost if 88 fully loaded 747 airliners crashed.”<sup>25</sup>

With regard to cerebral palsy, Calhoun et al 2007 calculated an estimated 1,096 cases of cerebral palsy each year attributable to induced surgical abortion and very preterm birth.<sup>23</sup>

Effects of abortion are not just neonatal: Gissler et al. 2004 found that pregnancy-related maternal mortality was three times as high for women within one year of abortion, compared to women after a live birth (83.1/100,000 compared to 28.2/100,000).<sup>27</sup> While this is likely related to many factors, it is important not to forget the maternal patient when thinking about the effects of abortion.

*Q What are the physician’s ethical obligations regarding this information?*

Ethical medical care requires informing women of the most recent and compelling evidence regarding the increased risk of subsequent PTB after a surgical abortion.

Informed consent remains a bedrock of ethical care for surgical and medical interventions. Patients deserve to know about of the risks associated with any procedure.

## Summary of Recommendations and Conclusion

*The following recommendations are based on good and consistent scientific evidence (Level A):*

1. Women who have a history of surgical abortion are at increased risk for preterm birth (delivery before 37 weeks).
2. Women who have a history of surgical abortion are at increased risk for very preterm birth (delivery before 32 weeks).
3. Multiple surgical abortions are associated with a “dose effect,” meaning more abortions confer more risk.

*The following recommendations are based on limited and inconsistent scientific evidence (Level B):*

1. Black Americans are disproportionately affected by abortion-related preterm birth.
2. The increased rate of preterm birth after surgical abortion is likely related to the surgical procedure itself.
3. There may be an inflammatory or subclinically infectious pathology associated with abortion-related preterm birth.
4. Women who have undergone medication abortions may be at increased risk

for preterm birth, especially if this was completed surgically.

*The following recommendations are based primarily on consensus and expert opinion (Level C):*

1. The relationship between abortion and preterm birth meets the Bradford Hill criteria for causality.
2. Abortion-related preterm birth has effects on neonates, mothers, and society at large.
3. Women with a previous history of termination of pregnancy should be informed of the increased risk for preterm birth.
4. Authors of studies and statements on preterm birth and abortion occasionally do not report their findings accurately.

## References

1. Lawn JE, Cousens S, Supan J; Lancet Neonatal Survival Steering Team. 4 million neonatal deaths: when? Where? Why? *Lancet* 2005;365(9462):891-900.
2. Tucker J, McGuire W. Epidemiology of preterm birth. *BMJ* 2004;329:675-678
3. Wen Sw, Smith G, Yang Q, Walker M. Epidemiology of preterm birth and neonatal outcome. *Semin Fetal Neonatal Med* 2004;9:429-435.
4. Martin JA., Kung HC, Mathews TJ, Hoyert DL, Strobine DM, Guyer B, Sutton SR. Annual summary of vital statistics: 2006. *Pediatrics* 2008;121:788-801.
5. Beck S. Wojdyloa D, Betran AP, Merialdi M, Requenjo JH, Rubens C, Menon R, Van Look



- PF. The worldwide incidence of preterm birth: a systematic review of maternal mortality and morbidity. *Bulletin World Health Organ* 2010;88:31-38.
6. Martin JA, Hamilton BE, Sutton PD, Ventura SJ, Menacker F, Munson ML. Births: Final data for 2002. *National Vital Statistics Reports*; Vol 52, No 10. Hyattsville, MD: National Center for Health Statistics; 2003.
  7. Centers for Disease Control and Prevention. Preterm Birth, (Nov 2021). Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention. <https://www.cdc.gov/reproductivehealth/maternalinfanthealth/pretermbirth.htm>
  8. Spong CY, Mercer BM, D'Alton M, Kilpatrick S, Blackwell S, Saade G. Timing of indicated late-preterm and early-term birth. *Obstet Gynecol* 2011;118:323-333.
  9. National Academies of Sciences, Engineering, and Medicine. 2018. *The Safety and Quality of Abortion Care in the United States*. Washington, DC: The National Academies Press. <https://doi.org/10.17226/24950>.
  10. Woolner A, Bhattacharya S, Battacharya S. The effect of method and gestational age at termination of pregnancy on future obstetric and perinatal outcomes: a register-based cohort study in Aberdeen Scotland. *BJOG* 2014;121:309-318.
  11. Bhattacharya S, Lowit A, Bhattacharya S, Raja EA, Lee AJ, Mahmood T, Templeton A. Reproductive outcomes following induced abortion: a national register-based cohort study in Scotland. *BMJ Open* 2012;2:e000911. Doi:10.1136/bmjopen-2012-00091.
  12. Liao H, Weu Q, Duan L, Ge J, Zhou Y, Zeng W. Repeated medical abortions and the risk of preterm birth in the subsequent pregnancy. *Arch Gynecol Obstet* 2011;284:579-586.
  13. Jackson, J. E., W. A. Grobman, E. Haney, and H. Casele. 2007. Mid-trimester dilation and evacuation with laminaria does not increase the risk for severe subsequent pregnancy complications. *International Journal of Gynaecology & Obstetrics* 96(1):12–15.
  14. Mirmilstein V, Rowlands S, King JF. *Aust N Z J Obstet Gynaecol*. 2009 Apr;49(2):195-7. doi: 10.1111/j.1479-828X.2009.00977.x
  15. Mannisto J, Bloigu A, Mentula M, Gissler M, Heikinheimo O, Niinimaki M. Interpregnancy interval after termination of pregnancy and risks of adverse outcomes in subsequent pregnancy. *Obstet Gynecol* 2017;129:347-54.
  16. Ball SJ, Pereira G, Jacoby P, de Klerk N, Stanley FJ. Re-evaluation of link between interpregnancy interval and adverse birth outcomes: retrospective cohort study matching two intervals per mother. *BMJ*. 2014 Jul 23;349:g4333. doi: 10.1136/bmj.g4333.
  17. Klebanoff MA. Interpregnancy Interval and Pregnancy Outcomes: Causal or Not? *Obstet Gynecol*. 2017 Mar;129(3):405-407. doi: 10.1097/AOG.0000000000001913.
  18. Interpregnancy Interval and Adverse Pregnancy Outcomes: An Analysis of Successive Pregnancies.
  19. Hanley GE, Hutcheon JA, Kinniburgh BA, Lee L. *Obstet Gynecol*. 2017 Mar;129(3):408-415. doi: 10.1097/AOG.0000000000001891.
  20. Thorp JM, Hartmann KE, Shadigian E. Long-term physical and psychological health consequences of induced abortion: review of



- the evidence. *Obstet Gynecol Survey* 2003;58(1):67-79.
21. Rooney B, Calhoun BC. Induced abortion and risk of later premature births. *J of Am Physicians and Surgeons* 2003;8(2):46-49.
  20. Iffy L. Letter. *Obstet Gynecol* 1975;45:115-116. Citing: Kovacs J: Nepesedespolitikank nehany kerdese: A kulong utodokert. *Magyar Hitek* 1973;26;10.
  21. Johnston WR. Historical abortion statistics, Hungary. Available at: <http://www.johnstonsarchive.net/policy/abortion/ab-hungary.html>.
  22. Calhoun BC, Shadigian E, Rooney B. Cost consequences of induced abortion as an attributable risk for preterm birth and impact on informed consent. *J Repro Med* 2007;52(10):929-937.
  23. McCaffrey M. The burden of abortion and the preterm birth crisis. *Issues in Law and Medicine* 2017;Vol 32, No 1:73-98.
  24. American College of Obstetricians and Gynecologists Compendium 2011. [www.acog.org](http://www.acog.org)
  25. Gissler et al 2004. Gissler M, Cerg B, Bouvier-Colle M, Bueens P. Pregnancy-associated mortality after birth, spontaneous abortion or induced abortion in Finland, 1987-2000. *Am J Ob Gyn* 2004;190:422-27.
  26. Oliver-Williams C, Fleming M, Monteath K., Wood AM, Smith GCS. Changes in Association between Previous Therapeutic Abortion and Preterm Birth in Scotland, 1980 to 2008: A Historical Chort Study. *PLOS* July 2013;10(7):1-11
  27. Jones RK and Jerman J. Abortion incidence an service availability in the Unites States, 2011 Perspective on Sexual and Reproductive Health 2014;46(1):3-14.
  28. Denison FC et al. The effect of mifepristone administration on leukocyte populations, matrix metallopreinases and inflammatory mediators in the first trimester cervix. *Mol Hum Reprod* 2000 Jun;6(6):541-548.
  29. Holt R. The molecular mechanisms of cervical ripening differ between term and preterm birth. *Endocrinology* 2001;15(3):1036-1046.
  30. Zhou Q, Zhang W, Xu H, Liang H, Ruan Y, Zhou S, LI X. Risk factors for preterm premature rupture of membranes in Chinese women from urban cities. *Int J Gynecology and Obstet* 2014;127:254-259.
  31. Lemmers M, Verschoor MA, Hooker AB, Opmeer BC, Limpens J, Huirne JA, Ankum WM, Mol BW. Dilatation and curettage increases the risk of subsequent preterm birth: a systematic review and meta-analysis. *Hum Repro* 2016 Jan;31(1):34-45.
  32. Hardy G, Benjamin A, Abenheim HA. Effect of induced abortions on early preterm births and adverse perinatal outcomes. *J Obstet Gynaecol Can* 2013;35(2):138-145.
  33. Martius JA, Steck T, Oehler MK, Wulf K-H. Risk factors associated with preterm (<37+0 weeks) and early preterm (<32+0 weeks): univariate and multi-variate analysis of 106 345 singleton births from 1994 statewide perinatal survey of Bavaria. *European J Obstetrics Gynecology Reproductive Biology* 1998;80:183-189.
  34. Maklouf MA, Clifton RG, Roberts JM, Myatt L, Hauth JC, Leveno KJ, Varner MW, Thorp JM Jr. Mercer BM, Peacemen AM, Ramin SM, Iams JD, Scioscione A, Tolosa JE, Sorokin Y. Eunice Kennedy Shriver National Institute of Child Health Human Development Maternal-Fetal Medicine Units Network. Adverse

- pregnancy outcomes among women with prior spontaneous or induced abortions. *Am J Perinatol* 2014 Oct;31(9):765-72.
35. Muhlemann K, Germain M, Krohn M. Does abortion increase the risk of intrapartum infection in the following pregnancy? *Epidemiology* 1996;7(2):194-198.
  36. Krohn M, Germain M, Muhlemann K. Prior pregnancy outcome and the risk of intra amniotic infection in the following pregnancy. *Am J Obstet Gyn* 1998;178(2):381-385.
  37. Hill AB. The environment and disease: association or causation? *Proc R Soc Med* 1965;58:295-300.
  38. Shah PS, Zao J. Induced termination of pregnancy and low birthweight and preterm birth: a systematic review and meta-analysis. *BJOG* 2009;116:1425-1442.
  39. Burguet A, Kaminski M, Abraham-Lerat L, Schaal JP, Cambonie G, Fresson J, Granjean H, Truffert P, Marpeau L, Voyer M, Roze JC, Treisser A, Larrroque B; EPIPAGE Study Group, *BJOG* 200411(3):258-265.
  40. Iams JD and Berghella V. Care for women with prior preterm birth. *Am J Obstet gynecol* 2011; 203(2):89-100.
  41. Steer P. (2009), Editors Choice *BJOG*; An International Journal of Obstetrics and Gynaecology 2009;116(11):i-ii.
  42. Royal College of Obstetrics and Gynecology "The Care of Women Requesting Induced Abortion." Evidence-based Clinical Guideline Number 7. Nov 2011. Available at [https://www.rcog.or.uk/globalassets/documents/guidelines/abortion-guide-line\\_wevb\\_1.pdf](https://www.rcog.or.uk/globalassets/documents/guidelines/abortion-guide-line_wevb_1.pdf) (last visited 30 May 18).
  43. <http://www.plannedparenthood.org/health-info/pregnancy/pregnant-now-what/thinking-about-abortion> (last visited 30 May 2018)
  44. Swingle HM, Colaizy TT, Zimmerman MB, Morriss FH. Abortion and the risk of subsequent preterm birth: A systematic review with meta-analyses. *J Rrepro Med* 2009;54:95-108.
  45. Oppenraaij RHF, Jauniaux E, Christianse OB, Horcajadas JA, Farquharson RG, Exalto N. Predicting adverse obstetric outcome after early pregnancy events and complications: a review. *Human Repro Update* 2009; 15(4):409-421.
  46. Lowit A, Bhattacharya S, Bhattacharya S. Obstetric performance following an induced abortion. *Best Practice and Research Clinical Obstetrics and Gynaecology* 2010; 24:667-682.
  47. Saccone G, Perriera L, Berghella V. Prior uterine evacuation of pregnancy as independent risk factor for preterm birth: a systematic review and metaanalysis. *Am J Obstet Gynecol* 2016;214:572-91.